

The Ly α Forest Flux Distribution at $z \sim 5.2$ and the Evolution of the Ionizing Background

Patrick McDonald^{1,2} and Jordi Miralda-Escudé^{1,3}

ABSTRACT

We use the redshift evolution of the Ly α forest mean transmitted flux at $z \gtrsim 2.5$ to infer the evolution of the intensity of the ionizing background, using theoretical predictions for the density distribution of the intergalactic gas giving rise to the Ly α forest. The proper background intensity declines gradually with redshift, decreasing by a factor ~ 3 between $z = 3$ and $z = 5$.

The gravitational structure formation theory of the Ly α forest implies that, given the observed mean transmitted flux of $\bar{F} = 0.1$ in the highest redshift quasar (with a mean Ly α forest redshift $z = 5.2$), only about 3% of the Ly α spectrum should have a flux higher than 50% of the continuum, and less than 0.5% of pixels should have flux above 75% of the continuum, assuming a uniform ionizing background intensity. We show that this is consistent with the spectrum shown in Fan et al. .

Subject headings: cosmology: theory—intergalactic medium—large-scale structure of universe—quasars: absorption lines

1. INTRODUCTION

The theory of the Ly α forest based on structure formation in Cold Dark Matter models has been highly successful in accounting for the observable properties of the Ly α forest, in particular the flux distribution and the power spectrum (see Croft et al. 1999, McDonald et al. 2000a, and references therein). As new quasars at ever higher redshift are discovered, the comparison of these observable quantities with the theory should be extended. Because the theory predicts the evolution of the spatial distribution of intergalactic gas, it is now possible to infer the evolution of the ionizing background intensity from the observed evolution of the mean flux decrement. Furthermore, as the

¹Department of Astronomy, The Ohio State University, Columbus, OH 43210; mcdonald,jordi@astronomy.ohio-state.edu

²Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104

³Alfred P. Sloan Fellow

epoch of reionization is approached, fluctuations in the background intensity due to the discreteness of the sources should increase, affecting the Ly α forest flux distribution.

In this paper, we use the observations of three quasars at $z \geq 5$ (Songaila et al. 1999, hereafter SHCM; Stern et al. 2000; Fan et al. 2000) to infer the evolution in the hydrogen ionization rate, and to test if the flux distribution is consistent with theoretical predictions. In §2 we present predictions of the flux distribution at high redshift, using numerical simulations whose resolution and box size are shown to be sufficient to have produced convergent results. In §3 we compare the predictions to the observations to measure the evolution of the ionization rate and search for differences in the predicted and observed flux distribution.

2. NUMERICAL RESULTS FOR THE FLUX DISTRIBUTION

If we ignore the effects of peculiar velocity of the gas and of thermal broadening, then the optical depth to Ly α scattering, τ , at any given point of an observed spectrum is determined by the density of gas at the corresponding point in space, $\rho_{\text{gas}} \equiv \Delta (\Omega_b \rho_{\text{crit}})$, and by the gas temperature, which we assume to follow the relation $T = T_0 \Delta^{\gamma-1}$ (see Hui & Gnedin 1997):

$$\tau = \tau_0 \frac{(1+z)^6 (\Omega_b h^2)^2}{T_0^{0.7} H(z) \Gamma_{-12}(z)} \Delta^\beta. \quad (1)$$

Here, τ_0 is a constant, $H(z) \simeq H_0 \Omega_m^{1/2} (1+z)^{3/2}$ is the Hubble constant at redshift z , H_0 is the present Hubble constant, $h = H_0/(100 \text{ km s}^{-1}/\text{Mpc})$, Ω_m is the fraction of the critical density in pressureless matter, $\Gamma_{-12}(z)$ is the photoionization rate of hydrogen in units of 10^{-12} s^{-1} , and $\beta = 2-0.7(\gamma-1)$ (where we use $\alpha(T) \propto T^{-0.7}$ for the recombination coefficient). The dependence of the distribution and average of the transmitted flux, $F = \exp(-\tau)$ (which is the observed quantity), on the cosmological model can be understood from equation (1). Thus, as long as the distribution of Δ and the $\Delta - T$ relation are sufficiently well understood, the observed mean transmitted flux of the Ly α forest can be used to determine the parameter combination $(\Omega_b h^2)^2/H(z)/\Gamma_{-12}(z)$ in equation (1).

Knowledge of the distribution of Δ and T needs to come from the theory, with the use of hydrodynamic simulations. This theory can be tested by checking its predictions for the detailed form of the distribution of F and its power spectrum against the observations, and this comparison has shown that the Cold Dark Matter model with a cosmological constant that is supported by a large body of observational evidence in galaxy clustering and CMB fluctuations (Primack 2000, and references therein) is in good agreement with the observed Ly α forest (Rauch et al. 1997, Croft et al. 1999, McDonald et al. 2000a, and references therein). At the same time, the gas temperature over the relevant range of densities has been measured from the absorption line shapes (Ricotti, Gnedin, & Shull 2000; Schaye et al. 2000; McDonald et al. 2000b), although the systematic differences between these studies are still substantial.

We focus in this paper on the flux distribution at high redshift. This particular property of the Ly α forest is affected only by the low-density intergalactic gas, given the large mean flux decrement (the reason is that high density gas causes absorption only in highly saturated regions of the spectrum). We first show that this distribution is predicted without significant uncertainties due to the finite resolution and box size of the simulations. We use the Hydro-PM (HPM) code of Gnedin & Hui (1998), comparing it first to the fully hydrodynamic simulation in Miralda-Escudé et al. (1996) to test that the approximate nature of the Hydro-PM method does not result in any significant differences in the predicted flux distribution at high redshift. When we create simulated spectra, we assume a $\Delta - T$ relation given by $T_0 = 2 \times 10^4$ K and $\gamma - 1 = 0$, which is consistent with the observational determinations Ricotti et al. (1999) and McDonald et al. (2000b) at $z \sim 4$, although Schaye et al. (2000) find a lower temperature (for the calculation of the pressure term in the HPM simulations we assume $T = 0$ before reionization at $z = 7$, and then interpolate between $T_0 = 25000$, $\gamma - 1 = 0.0$ at $z = 7$ and $T_0 = 19000$, $\gamma - 1 = 0.2$ at $z = 3.9$; however, we find that the results are almost completely insensitive to these assumptions, probably because the pressure does not have much influence in the voids). The cosmological model is Λ CDM, with $\Omega_m = 0.4$, $\Omega_b h^2 = 0.015$ (note, however, that the results we give for Γ_{-12} are rescaled to $\Omega_b h^2 = 0.02$), $\Omega_\Lambda = 0.6$, $h = 0.65$, and $n = 0.95$, where Ω_Λ is the fraction of the critical density in cosmological constant and n is the usual logarithmic slope of the primordial power spectrum. Simulated spectra are constructed in which the transmitted flux is binned into $2 \text{ \AA}$ pixels, to match the observed spectrum in Fan et al. (2000) (we use this pixel size throughout the paper). In Figure 1, we show the flux distribution predicted for the full-hydro simulation as the heavy solid line. The light (*long-dashed*, *dotted*, *solid*, *dashed*) lines show HPM simulations with box size $L = (17.8, 8.9, 4.4, 4.4)$, and a total number of particles $N = (256^3, 128^3, 128^3, 256^3)$, respectively. All the results are at $z = 4$, but using the value of Γ_{-12} that gives a mean flux decrement from the hydrodynamic simulation of $\bar{F} = 0.1$, which matches the observations at $z \sim 5$. We keep Γ_{-12} (rather than $\bar{F}$) constant for all the simulations, so that we can see the difference in the predicted $\bar{F}$.

The result for the fully hydrodynamic simulation and for the HPM simulation that has almost the same box size and resolution ($L = 8.9$ and $N = 256^3$, not plotted) are essentially identical. This justifies using the HPM simulations to examine the effects of changing the box size and resolution and to compute the flux distribution for comparison with observations at different redshifts. The two simulations with the smallest box size, with resolutions $L/N^{1/3} = 35 h^{-1} \text{ Kpc}$ and $17 h^{-1} \text{ Kpc}$ (solid and dashed lines, respectively), show that simulations with inter-particle separation of $35 h^{-1} \text{ Kpc}$ have already achieved the required resolution; however, degrading the resolution to $L/N^{1/3} = 69 h^{-1} \text{ Kpc}$ causes a change in $\bar{F}$ of 0.02, similar to the observational error bars of $\bar{F}$. Comparison of the long-dashed and dotted (or the heavy and light solid) lines shows that the effect of increasing the box size beyond $5 h^{-1} \text{ Mpc}$ is small. We conclude that simulations with boxes $L = 8.9 h^{-1} \text{ Mpc}$ and number of particles $N = 256^3$ are sufficient for our purpose to have achieved convergence. We find similar convergence in the HPM simulations at $z = 5.2$. We also find that different random realizations of the initial conditions of the simulation of this box size and resolution change $\bar{F}$ by ± 0.01 (small compared to the observational error).




Fig. 1.— Cumulative distribution of transmitted flux. *Heavy solid line*: full hydrodynamic simulation. *Thin lines*: Hydro-PM simulations, with box sizes and particle numbers indicated.

3. RESULTS

We now examine the flux distribution at high redshift predicted from the simulations to do a direct comparison to the observational constraints. We use the HPM simulation with $L = 8.9 h^{-1} \text{ Mpc}$ and $N = 256^3$, and create spectra binned into $2 \text{ \AA}$ pixels. The curves in Figure 2(a) show the cumulative flux distribution at $z = 5.2$, for ten values of Γ_{-12} , corresponding to different values of $\bar{F}$, indicated in the Figure. This can be compared to the spectrum in Figure 3 of Fan et al. (2000), from a quasar at $z = 5.8$ (where the mean redshift of the observed Ly α forest, in the wavelength range 7140-7956 $\text{\AA}$ used by Fan et al. 2000 to compute $\bar{F}$, is 5.2). In order to match the observed $\bar{F} = 0.10 \pm 0.02$, we need $\Gamma_{-12} \simeq 0.17 \pm 0.03$, for the Ly α forest model described in §2.

Remarkably, there should be essentially no pixels in the spectrum at this high redshift where the flux comes close to the continuum level. Only 3% of the spectrum should have a flux higher than 50% of the continuum, and less than 0.5% should have a flux greater than 75% of the continuum [see the inset of Fig. 2(a)]. In comparison, Fan et al. (2000) identify the minimum absorption in their observed spectrum to be $\tau \sim 0.4$ (although they estimate that continuum uncertainty and noise make this number uncertain by about ± 0.1). Absorption of $\tau \lesssim 0.4$ or $F \lesssim 0.67$ occurs in approximately 1% of the pixels in our simulated spectra, so we predict about three such pixels to be found in the wavelength range 7300-7956 $\text{\AA}$ examined by Fan et al. (2000). Therefore, the minimum absorption present in this spectrum is consistent with the predictions of the simple Ly α forest model in our simulations. We find that decreasing σ_8 by $\sim 15\%$ to match the results of McDonald et al. (2000a) increases the inferred value of Γ_{-12} by $\sim 30\%$, but does not affect significantly our conclusions about the flux distribution (once the observed $\bar{F}$ is matched).

Next, the same lines are plotted in Figure 2(b) at redshift $z = 4.5$, which is about the mean redshift of the Ly α forest in the $z = 5.0$ quasar spectrum that is shown in Figure 2 of SHCM. The mean transmitted flux in this spectrum is $\bar{F} = 0.25$, which requires $\Gamma_{-12} \simeq 0.21$ (no error bar was given in SHCM for the value of $\bar{F}$). According to SHCM, an upper bound on the minimum optical depth in their spectrum is $\tau < 0.1$. They identify this region as being produced by a void with density $\Delta \simeq 0.2$, to derive a lower limit to the background intensity of $\Gamma_{-12} > 2.8$, for $\Omega_b h^2 = 0.019$.

Our value of Γ_{-12} , inferred by matching $\bar{F}$, is much smaller for the following reasons: Factors of $\sim (20000/5000)^{-0.7}$, $(0.4/1.0)^{1/2}$, and $0.65/0.5$ are obtained from our different choices of T_0 , Ω_m , and h , respectively (note: we assume SHCM used $h = 0.5$, but their paper is not clear on this point). Our choices place more conservative lower limits on Γ_{-12} , and are more consistent with independent observations. The remaining factor 2.4 must be traced to the identification by SHCM of observed $\tau = 0.1$ with underdensity $\Delta = 0.2$. In our simulations, $\Delta < 0.2$ occurs in 1.7% of pixels in real space, and this probability is increased when the expansion of voids in redshift space is considered. Consequently, the minimum absorption in the $200 \text{\AA}$ region considered by SHCM probably arises from a void with $\Delta < 0.2$. Furthermore, the peculiar expansion of voids reduces the optical depth at fixed density. After the inclusion of all these effects through our simulations, and using $\Gamma_{-12} = 0.21$,





Fig. 2.— Cumulative flux distributions for (a) $z = 5.2$ and (b) $z = 4.5$. Listed values of Γ_{-12} and $\bar{F}$ apply to the lines from top to bottom. The inset box in (a) is an expanded version of the upper right hand corner of the full figure.



we predict ~ 0.4 regions with $\tau < 0.1$ in the wavelength range $6800 - 7000 \text{ \AA}$ identified by SHCM. Lastly, the minimum absorption statistic is biased toward regions with strong noise contribution, so the value $\tau < 0.1$ is probably not conservative (our simulations give a completely acceptable 1% probability for $\tau < 0.15$).

Finally, the quasar in Stern et al. (2000) at $z = 5.5$ has mean transmitted flux $\bar{F} = 0.10 \pm 0.02$. The ionization rate required to match this $\bar{F}$ is $\Gamma_{-12} = 0.13 \pm 0.03$, at a mean redshift $\bar{z} = 4.93$. Their spectrum as shown has only $\sim 15 \text{ \AA}$ resolution so we do not interpret it further.

The evolution of the background intensity Γ_{-12} derived from the observed mean flux decrement, for $\Omega_b h^2 = 0.02$, is shown in Figure 3, using the values from the three quasars discussed here and the lower redshift results from McDonald et al. (2000a) (updated by McDonald et al. 2000b using the measured IGM temperatures).

4. DISCUSSION

There is a clear reduction of the proper intensity of the ionizing background with redshift, by a factor ~ 3 from $z = 3$ to $z = 5$. This reduction was noticed previously by Rauch et al. (1997) up to $z = 4$. With the new high redshift quasars, we find a continuation of this gradual decrease of Γ , but without a dramatic drop that would be the sign of approaching the reionization epoch. The consistency of the fraction of the Ly α spectrum with low absorption with the observations also suggests that the background intensity fluctuations are not yet extreme at $z = 5$.

The evolution of the background intensity is related to the evolution of the comoving emissivity, ϵ , and the proper mean free path λ , by $\epsilon (1+z)^3 \propto \Gamma \lambda^{-1}$, if λ is small compared to the cosmic horizon. Assuming that the rate of incidence of Lyman limit systems evolves as $(1+z)^{1.5}$ (Storrie-Lombardi et al. 1994), the mean free path of ionizing photons is $\lambda \propto (1+z)^{-4}$, and therefore the comoving emissivity changes between $z = 3$ and $z = 5$ by a factor $(1/3)(6/4)^{4-3} = 0.5$, indicating only a small reduction, which is clearly different from the reduction by a factor 0.14 predicted by the evolution of the comoving density of luminous quasars (Schmidt, Schneider, & Gunn 1995) and implies, therefore, that galaxies or other sources are dominating the emission of ionizing photons at $z = 5$.

We thank Nick Gnedin for providing a copy of his Hydro-PM code and for helpful comments on the manuscript, Renyue Cen for the output from his hydrodynamic simulation, and Adam Steed for helpful comments on the manuscript.

REFERENCES

- Croft, R. A. C., Weinberg, D. H., Pettini, M., Hernquist, L., & Katz, N. 1999, ApJ, 520, 1
 Fan, X. et al. 2000, AJ, submitted (astro-ph/0005414)




Fig. 3.— Evolution of observed Γ_{-12} . Computed for $\Omega_B h^2 = 0.02$, and the Λ CDM model discussed in the text (changing Ω_B , for example, will shift all the points by the same factor). Uncertainties related to the quasar continuum and the gas temperature may introduce some uncertainty beyond the given error bars, especially for the 3 highest z points.

- Gnedin, N. Y. & Hui, L. 1998, MNRAS, 296, 44
- Hui, L. & Gnedin, N. Y. 1997, MNRAS, 292, 27
- McDonald, P., Miralda-Escudé, J., Rauch, M., Sargent, W. L. W., Barlow, T. A., Cen, R., & Ostriker, J. P. 2000a, ApJ, in press (astro-ph/9911196)
- McDonald, P., Miralda-Escudé, J., Rauch, M., Sargent, W. L. W., Barlow, T. A., & Cen, R. 2000b, ApJ, submitted (astro-ph/0005553)
- Miralda-Escudé, J., Cen, R., Ostriker, J. P., & Rauch, M. 1996, ApJ, 471, 582
- Primack, J. R. 2000, preprint (astro-ph/0007187)
- Rauch, M., Miralda-Escudé, J., Sargent, W. L. W., Barlow, T. A., Weinberg, D. H., Hernquist, L., Katz, N., Cen, R., & Ostriker, J. P. 1997, ApJ, 489, 7
- Ricotti, M., Gnedin, N. Y., & Shull, J. M. 2000, ApJ, 534, 41
- Schaye, J., Theuns, T., Rauch, M., Efstathiou, G., & Sargent, W. L. W. 2000, MNRAS, in press (astro-ph/9912432v2)
- Schmidt, M., Schneider, D. P., & Gunn, J. E. 1995, AJ, 110, 68
- Songaila, A., Hu, E. M., Cowie, L. L., & McMahon, R. G. 1999, ApJ, 525, L5 (**SHCM**)
- Stern, D., Spinrad, H., Eisenhardt, P., Bunker, A. J., Dawson, S., Stanford, S. A., & Elston, R. 2000, ApJ, 533, L75
- Storrie-Lombardi, L. J., McMahon, R. G., Irwin, M. J., & Hazard, C. 1994, ApJ, 427, L13